# EXOTHOR

## INSECT GROWTH REGULATOR

Registration No. L11062 of Act No. 36 of 1947 Namibian Reg. No. N-AR 2264

A water dispersible granule for the prevention of the development of immature cockroaches, flies, fleas and mosquitoes to adulthood.

**ACTIVE INGREDIENT:** 

200 g/kg DIFLUBENZURON (benzoylurea)

**Net Contents:** 

500 g

## **ENSYSTEX SOUTH AFRICA (PTY) LTD**

Building 3, The Burns Office Park 12 A Jet Park Road, Jet Park Boksburg, 1469 www.Ensystex.co.za

Customer Service Call 080 367 978

®™ Trademarks of Ensystex, Inc. used by licence to Ensystex South Africa (Pty) Ltd.





CAUTION VERSIGTIG





**ENSYSTEX** 

## DIRECTIONS FOR USE: USE ONLY AS DIRECTED

SITUATION	PEST	Dosage rate	Remarks
Fly breeding areas, e.g. manure and rubbish dumps, at domestic, commercial, industrial and public buildings and facilities.	House fly maggots Musca domestica	75 g / 10 L of water / 20 m <sup>2</sup> 37.5 g / 10 L of water / 20 m <sup>2</sup>	Apply to fly breeding areas at a rate of 500 mL per m². Use the higher rate for best results or where fly populations are larger. Use the lower rate for maintenance treatments or when fly populations are smaller.
Domestic, commercial, industrial and public buildings and facilities.	Cockroach nymphs Blattella germanica and Periplaneta americana	75 g / 10 L of water / 100 $\mbox{m}^2$	Apply as a coarse spray at the rate of 1 litre mixture per 10 m², or to the point of run-off which will vary with the nature of the surface. Reapply when signs of infestation reappear. Apply to areas where insects harbour, paying particular attention to cracks and crevices.
	Fleas (immature stages)	150 g / 10 L of water / 100 m <sup>2</sup>	Apply as a course spray at the rate of 1 litre mixture per 10 m <sup>2</sup> , to carpets. Reapply when signs of infestation reappear.
Mosquito breeding sites	Mosquito larvae of the <i>Anopheles</i> , <i>Aede</i> s and <i>Culex</i> genera	Clear water: 250 - 500 g / ha (2.5 - 5.0 g / 100 m²) Polluted water: 500 - 1000 g / ha (5.0 - 10 g / 100 m²)	Apply at an early stage of population development when the majority of the larvae are in the first or second instar.  Applications should be repeated when inspection indicates an increase in the number of second and third instar of the mosquito larvae.  Apply in sufficient water to ensure even coverage of breeding sites.

### WARNING

Handle with Care.

May cause slight eye irritation.

Toxic to crustaceans and other aquatic and soil arthropods. Keep out of reach of children, uninformed persons and animals.

### **PRECAUTIONS**

Do not inhale the spray mist and avoid contact with eyes. Wear protective clothing during preparation of spray mixture

- face shield and protective gloves.

Wash with soap and water after use or accidental skin contact. Wash contaminated clothing after a day's work.

Do not eat, drink or smoke whilst mixing or applying and before washing hands and face.

Clean applicator thoroughly after use and do not leave spray mixture standing in applicators overnight.

Dispose of wash water where it will not contaminate food, grazing or water.

Triple rinse the empty container as follows: invert the container over the spray or mixing tank and empty thoroughly.

Thereafter, rinse the container three times with a volume of water equal to a minimum of 10% of that of the container. Add the rinsings to the contents of the spray tank before destroying the container.

Do not use the empty container for any other purpose.

Do not apply this product to animals.

Remove pets and cover or remove aquaria from the area to be treated.

Do not walk across treated areas (e.g. carpets) until application has dried thoroughly.

Avoid contamination of food, feed drinking water and eating utensils.

## NOTICE

Ensystex warrants that this product conforms to its chemical description and is reasonably fit for the purposes stated on the label when used in accordance with Directions for Use under normal conditions of use. No warranty of merchantability or fitness for a particular purpose, express or implied, extends to the use of the product contrary to label instructions or under abnormal conditions.

Date of Manufacture:

9 330209 005065 >

L-SA-CHEEX1004 1.02 11.22