



RODENTHOR[®]

PASTA

Reg. No. L9586 Act/Wet No. 36 of/van 1947
Namibian Reg. No. N-AR1647

A ready-for-use pasta bait for the control of roof and Norway rats and the house mouse in and around industrial premises and human dwellings.

'n Gereed-vir-gebruik pasta-lokaas vir die beheer van dak - en Noorweegse rotte en die huismuis in en om industriële persele en wonings van mense.

ACTIVE INGREDIENT / AKTIEWE BESTANDDEEL

Brodifacoum (coumarin anticoagulant).....0,05 g/kg

Brodifakum (kumarinbloedstolweermiddel).....0,05 g/kg

Batch No:
Lotnr:

Date of manufacture:
Datum van vervaardiging:

SOUTH AFRICA REGISTRATION HOLDER /
REGISTRASIEHOUER:
Mr. Stefano Migliore
TEL.: 011-453-5638

DISTRIBUTOR / VERSPREIDER:
ENSYSTEX SOUTH AFRICA (PTY) LTD
Co./Mpy Reg. No. 2008/009219/07
Building 3, The Burns Business Park
12A Jet Park Road, Jet Park
Boksburg GAUTENG 1469
Tel.: 080-367-9783 www.Ensystex.co.za

NET MASS /
NETTO MASSA **3 kg**

UN number/VN-nommer: 3808

ENSYSTEX[®]



HARMFUL
SKADELIK



WARNINGS

- Handle with care.
- Harmful when swallowed and by skin contact.
- A mild skin and eye irritant.
- Toxic to fish and other animals.
- Store in a cool place away from food and feedstuffs.
- Keep out of reach of children, uninformed persons and animals.
- In case of poisoning call a doctor and show him this label.

PRECAUTIONS

- Avoid contact with skin.
- Wear protective gloves when handling bait.
- After use and in case of accidental skin contact, wash thoroughly with soap and water.
- Destroy empty container and do not re-use for any other purpose.
- Prevent contamination of food, feedstuffs, eating utensils and drinking water.
- Set bait where it will be inaccessible to children and non target animals.

SYMPTOMS OF HUMAN POISONING

Bleeding from gums and nose, bleeding under the skin, pallor, haemorrhage into and around elbow and knee joints. Blood in urine and faeces.

FIRST AID TREATMENT

Eyes – Rinse eyes with plenty of clean water for at least 15 minutes.

Skin – Wash contaminated body area thoroughly with plenty of soap and water.

Ingestion – Induce vomiting by tickling back of throat. Call a doctor and show him this label.

NOTE TO PHYSICIAN

Administer Vitamin K intravenously, 50-150 mg depending on prothrombin time on first day. In severe cases small transfusions of fresh blood can be considered.

DIRECTIONS FOR USE

Use only as indicated.

RODENTHOR PASTA kills after rodents have had one or more feeds. Death will occur 4-12 days after a lethal dose has been consumed.

Each sachet contains 10 g of **RODENTHOR PASTA**.

BAITING

1. Place **RODENTHOR PASTA** bait at a suitable number of locations near where rodents have nests or burrows or where they have been feeding or caused damage. Inspect bait point regularly and top-up until rodent activity ceases.
2. Dosage rates:
Rats: 50-100 g **RODENTHOR PASTA** at a baiting point at intervals of 5-10 meters.
Mice: 20-30 g **RODENTHOR PASTA** at a baiting point at intervals of 2-5 meters.
3. Wear PVC or rubber gloves when handling bait as rodents are deterred by human odour.
4. Ensure that domestic and farm animals do not have access to bait by placing in a piece of drain pipe or under a tile.
5. Replace completely any bait which has become spoilt or contaminated and dispose of in a responsible manner.
6. When rodent activity has ceased, remove uneaten bait and destroy in a responsible manner. It is good practice however, to maintain a few permanent feeding points to prevent reinfestation.

WAARSKUWINGS

- Hanteer versigtig.
- Skadelik indien ingesluk en in aanraking met vel.
- 'n Ligte vel- en oogirritant.
- Giftig vir vis en ander diere.
- Berg in 'n koel plek weg van voedsel en voer.
- Hou buite die bereik van kinders, oningeligte persone en diere.
- In geval van vergiftiging ontbied 'n geneesheer en toon hierdie etiket aan hom.

VOORSORGMATREËLS

- Voorkom aanraking met vel.
- Dra beskermende handskoene wanneer lokaas hanteer word.
- Na gebruik en in die geval van toevallige aanraking met vel was deeglik met seep en water.
- Vernietig leë houers en moet nie vir enige ander doel hergebruik nie.
- Voorkom besoedeling van voedsel, voer, eetgerei en drinkwater.
- Plaas lokaas sodat dit ontoeganklik vir kinders en nie-teiken diere sal wees.

SIMPTOME VAN MENSLIKE VERGIFTIGING

Bloeding van die tandvleis en neus, bloeding onder die vel, bleekheid, bloeding in en om elmboog- en kniegewrigte, bloed in die uriene en ontlasting.

NOODHULPBEHANDELING

Oë – Spoel oë vir minstens 15 minute met volop skoon water.

Vel – Spoel besoedelde liggaamsdele met volop seep en water.

Inname – Bring braking te weeg deur agterkant van keel te prikkel. Ontbied 'n geneesheer en stel hierdie etiket aan hom beskikbaar.

NOTA AAN GENEESHEER

Dien Vitamien K binne-aars toe teen 50-150 mg afhange van protrombientyd op die eerste dag. In ernstige gevalle kan klein bloedtransfusies met vars bloed oorweeg word.

GEBRUIKSAANWYSINGS

Gebruik slegs soos aangedui.

RODENTHOR PASTA dood rotte en muise na een of meer innames van die lokaas. Knaagdiere sal 4-12 dae neem om dood te gaan nadat 'n dodelike dosis ingeneem is.

Elke sakkie (sachet) bevat 10 g **RODENTHOR PASTA**.

PLASING VAN LOKAAS

1. Plaas **RODENTHOR PASTA** by 'n voldoende aantal plekke naby muis- en rotgate of neste, of waar die knaagdiere vreet of skade aanrig, uit. Ondersoek die lokaaspunte gereeld en vul aan met lokaas totdat rot- en muisaktiwiteit ophou.
2. Dosisse:
Rotte: 50-100 g **RODENTHOR PASTA** per lokaaspunt 5-10 meters uitmekaar.
Muise: 20-30 g **RODENTHOR PASTA** per lokaaspunt 2-5 meters uitmekaar.
3. Dra PVC- of rubberhandskoene wanneer lokaas hanteer word omdat knaagdiere afgeskrik word deur die reuke van mense.
4. Dra sorg dat huis- en plaasdiere nie toegang tot lokaas het nie deur bv. dit in 'n stuk rioolpyp of onder 'n teël te plaas.
5. Vervang alle bedorwe of gekontamineerde lokaas en raak daarvan ontslae op 'n verantwoordelike wyse.
6. Wanneer knaagdiereaktiwiteit ophou, verwyder en vernietig oorblywende lokaas op 'n verantwoordelike wyse. Dit is egter raadsaam om 'n paar permanente lokaaspunt te behou ten einde herbesmetting te voorkom.

